

Genesis Biotech Inc.

http://www.genesisbio.com.tw info@genesisbio.com.tw

TEL: +886-2-22181731 FAX: +886-2-22181732 Date: 02/17/2011

Anti-Estrogen receptor 2 (ESR2), Chicken-Polyclonal Antibody

Catalog No. PY-10160 Antigen species: Human Host species: Chicken Quantity:100μgApplications tested:Western blotReactivity:Human, Mouse, RatForm:Antigen affinity purified antibody

Target description

Estrogen receptor-beta (ESR2) is a member of the superfamily of nuclear receptors, which can transduce extracellular signals into transcriptional responses.

Antigen

This polyclonal antibody was raised by immunizing chicken with ESR2 fusion protein (1-150 amino acids).

Application

Western blotting, tissue or cell immunostaining. Recommended starting dilution for Western blot analysis is 1:500, for tissue or cell staining is 1:200. Optimal working dilutions must be determined by the end user.

Related Products

- 1. Anti-Cytokeratin 8 (KRT8), Chicken pAb (PY-10149).
- 2. Anti-Cytokeratin 18 (KRT18), Chicken pAb (PY-10150).
- 3. Anti- Toll-like receptor 2 (TLR2), Chicken pAb (PY-10152)
- 4. Anti- Toll-like receptor 3 (TLR3), Chicken pAb (PY-10154)
- 5. Anti- Toll-like receptor 7 (TLR7), Chicken pAb (PY-10148)



E coli-derived fusion protein as test antigen.

Western blot Protocol

- 1. Block membrane with 5% non-fat milk in PBS-T for 1 hour at room temperature or longer at 4°C.
- 2.Incubate membrane with IgY antibodies at dilution of 1: 2,000 with 1% milk in PBS-T at R.T. for 1 h.
- 3.Rinse 3 times with PBS-T, then wash membrane with PBS-T, 5 min each, total of 3 times.
- 4.Incubate with 2nd antibody (goat-anti-IgY/Fc-HRP) at dilution 1:1,000 for ECL (with 1%milk PBS-T) at R.T. for 1h.
- 5.Rinse 3 times with PBS-T, then wash with PBS-T, 5 min each with shaking, total of 3 times.
- 6.Perform ECL detection of signal using Pierce ECL kit.

Storage

It is supplied as antigen affinity purified antibody in lyophilized powder. Redissolve the powder with 100 microliter sterile water will restore to the original concentration 1mg/ml (1×PBS). Store at 4° C for short-term application. For longterm storage, aliquot and store at -20°C.

References

Weihua, Z.; Saji, S.; Makinen, S.; Cheng, G.; Jensen, E. V.; Warner, M.; Gustafsson, J.-A. : Estrogen receptor (ER)-beta, a modulator of ER-alpha in the uterus. Proc. Nat. Acad. Sci. 97: 5936-5941, 2000.

FOR RESEARCH USE ONLY AND NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE