



Genesis Biotech Inc.

website: <http://www.genesisbio.com.tw> e-mail : info@genesisbio.com.tw

Anti-Beta-actin, Chicken polyclonal Antibody

Catalog No. PY-10190 **Quantity:** 100 µg **Host species:** chicken
Reactivity: Bovine, human, mouse, rat, rabbit, and many other species.
Applications tested: W. B. **Type:** polyclonal antibody **Form:** Affinity-purified IgY Ab

Target description

Beta actin is one of six different actin isoforms which have been identified. ACTB is one of the two nonmuscle cytoskeletal actins. Actins are highly conserved proteins that are involved in cell motility, structure and integrity. Alpha actins are a major constituent of the contractile apparatus.



Antigen

This polyclonal antibody was raised by immunizing chicken with beta-actin fusion protein (151-320 amino acids).

293 cell lysate as test antigen.

Application

Western blotting, tissue or cell immunostaining. Recommended starting dilution for Western blot analysis is 1:1000, for tissue or cell staining is 1:200. Optimal working dilutions must be determined by the end user

Western blot Procedure

1. Block membrane with 5% non-fat milk in PBS-T for 1 hour at room temperature or longer at 4°C.
2. Incubate membrane with IgY antibodies at dilution of 1: 1,000 with 1% milk in PBS-T @ R.T. for 1 h.
3. Rinse 3 times with PBS-T, then wash membrane with PBS-T, 5 min each, total of 3 times.
4. Incubate with 2nd antibody (goat-anti-IgY/Fc-HRP) at dilution 1:1,000 for ECL (with 1% milk PBS-T) @ R.T. for 1 h.
5. Rinse 3 times with PBS-T, then wash with PBS-T, 5 min each with shaking, total of 3 times.
6. Perform ECL detection of signal using Pierce ECL kit.

Related Products

Anti-Laminin, Chicken-Poly (PY-10138)
Anti-Histone H1, chicken-poly (PY-10140)

Storage

It is supplied as affinity purified antibody/ PBS buffer in lyophilized powder. Redissolve the powder with 100 µL sterile water will restore to the original concentration 1mg/mL. Store at 4°C for short-term application. For long-term storage, aliquot and store at -20°C.

References

Lloyd, C. and Gunning, P. beta- and gamma-actin genes differ in their mechanisms of down-regulation during myogenesis J. Cell. Biochem. 84 (2), 335-342 (2002)

FOR RESEARCH USE ONLY AND NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE